


4. (Amended) Epothilone derivative of the formula (5)



wherein the residue R^1 is a hydrogen atom or a C_{1-8} -alkyl group, and P is a protective group and X-Y is a group of formula $-CH_2CH-OP$ or $CH=CH$, and R^2 is a monocyclic aromatic which can be substituted by a halogen atoms and/or OR^4 - and/or NR^5R^6 - and/or alkyl, alkenyl and/or alkynyl groups in ortho- and/or meta- and/or para-position, or a monocyclic 5- or 6-membered hetero aromatic, which can be provided with one or several O- and/or N- and/or S-atoms in the ring and/or which can be provided with OR^4 - and/or NR^5R^6 - and/or alkyl, alkenyl and/or alkynyl groups as substituents, wherein the residues R^4 , R^5 and R^6 independently are defined as R^1 in claim 1, but are independent of R^1 , wherein

(i) XY is excluded as group of formula $-CH=CH-$ if R^1 is a hydrogen atom or a C_{1-4} -alkyl group and R^2 is a monocyclic hetero aromatic having a N atom and/or a S atom in its ring and a C_1 -alkyl substituent and

(ii) XY is excluded as group of formula $-CH_2-CH-OP$ if R^1 is a hydrogen atom or a C_{1-4} -alkyl group and R^2 is a monocyclic hetero aromatic having a N atom and/or a S atom in its ring and a C_1 -alkyl substituent.

Swb
E3
com
T,0200


C1
com

~~6.~~ (Amended) Epothilone derivative as in claims 1, 2, 3, 4, 5 or ~~22~~ wherein R^1 , R^4 , R^5 and R^6 are a hydrogen atom or a C_{1-6} -alkyl group.

~~7.~~ (Amended) Epothilone derivative as in claims 4, 5, ~~6~~ or ~~22~~ wherein the substituents of the monocyclic aromatic and/or hetero aromatic are C_{1-6} -alkyl, C_{2-6} -alkenyl and C_{2-6} -alkinyl groups respectively, especially C_{1-4} -alkyl, C_{2-4} -alkenyl and C_{2-4} -alkinyl groups, respectively and fluoro, chloro, bromo or iodo atoms.


~~8.~~ (Amended) Epothilone derivative as in claims 4, 5, ~~6~~, ~~7~~ or ~~22~~ wherein the aromatic and hetero aromatic, respectively, is provided with 1, 2 or 3 substituents and the hetero aromatic is provided with 1, 2 or more hetero atoms.

~~9.~~ (Amended) Process for the preparation of a compound of formula (3),




wherein a compound of formula (2) according to claim 1 is reacted with the compound of formula $\text{HC} [\text{B} (\text{OR})_2]_3$ optionally in the presence of a base, wherein the residue R^1 is a hydrogen atom or a C_{1-8} -alkyl group, X-Y is a group of formula $-\text{CH}_2\text{CH}-\text{OP}$ or $-\text{CH}=\text{CH}-$, and P is a protective group, wherein X-Y is excluded as group of formula $-\text{CH}_2\text{CH}-\text{OP}$ if R^1 means a hydrogen atom or a C_{1-4} -alkyl group and R is defined as R^1 , but is independent of R^1 .

11/10. (Amended) Process for the preparation of a compound of formula (4),




wherein a compound of formula (3) according to claim 2 is reacted with N-iodo- or N-bromo succinimide and that the residue R^1 is a hydrogen atom or a C_{1-8} -alkyl group, X-Y is a group of formula $-\text{CH}_2\text{CH}-\text{OP}$ or $-\text{CH}=\text{CH}-$, and P is a protective group, wherein X-Y is excluded as group of formula $-\text{CH}_2\text{CH}-\text{OP}$ if R^1 means a hydrogen atom or a C_{1-4} -alkyl group.

12/11. (Amended) Process for the preparation of a compound of formula (5),



wherein a compound of formula (3) according to claim 2 is reacted by a Suzuki coupling with a compound of formula R^2-Z , wherein R^2 is a monocyclic aromatic which can be substituted by halogen atoms and/or OR^4 - and/or NR^5R^6 - and/or alkyl, alkenyl and/or alkynyl groups in ortho- and/or meta- and/or para-position, or a monocyclic 5- or 6-membered hetero aromatic, which can be provided with one or several O- and/or N- and/or S-atoms in the ring and/or which can be provided with OR^4 - and/or NR^5R^6 - and/or alkyl, alkenyl and/or alkynyl groups as substituents and Z can be a halogen atom or a group of formula $-OSO_2CF_3$, $-CH=CHI$, $-CH=CHOSO_2CF_3$.


13/ 12. (Amended) Process for the preparation of a compound of formula (5),



C₂
C₂


wherein a compound of formula (4) according to claim 3 is reacted by a silent coupling (stille Kupplung) with R_2-SNR^3 , wherein R^2 is a monocyclic aromatic which can be substituted by halogen atoms and/or OR^4 - and/or NR^5R^6 - and/or alkyl, alkenyl and/or alkynyl groups in ortho- and/or meta- and/or para-position, or a monocyclic 5- or 6-membered hetero aromatic, which can be provided with one or several O- and/or N- and/or S-atoms in the ring and/or which can be provided with OR^4 - and/or NR^5R^6 - and/or alkyl, alkenyl and/or alkynyl groups as substituents and R^3 is a C_{1-6} -alkyl group.

14 13. (Amended) Process for the preparation of a compound of formula (6),



wherein the protective group is removed from a compound of formula (5) according to claim 4.

15 14. (Amended) Process for the preparation of a compound of formula (6),




wherein it comprises the process steps as disclosed in claims 9, 10, 11, 12 or 13.

16 15. (New) A pharmaceutical composition comprising at least one of the compounds described in claims 1, 2, 3, 4, 5, 6, 7, 8 or 22 and optionally carriers, diluents and/or auxiliary agents.

18

22. (New) Epothilone derivative of formula (6)



wherein the residues are defined as in claim 4 and, if X-Y means a group of formula $-\text{CH}_2\text{CH}-\text{OP}$, the protective group P has been removed, wherein

(i) XY is excluded as group of formula $-\text{CH}=\text{CH}-$ if R^1 is a hydrogen atom or a C_{1-4} -alkyl group and R^2 is a monocyclic hetero aromatic having a N atom and/or a S atom in its ring and a C_1 -alkyl substituent and

(ii) XY is excluded as group of formula $-\text{CH}_2\text{CH}-\text{OP}$ if R^1 is a hydrogen atom or a C_{1-4} -alkyl group and R^2 is a monocyclic hetero aromatic having a N atom and/or a S atom and/or an O atom in its ring and a C_1 -alkyl substituent.

23. (Amended) Epothilone derivative according to claim 22, wherein the substituents of the monocyclic aromatic and/or hetero aromatic are C_{1-6} -alkyl, C_{2-6} -alkenyl and C_{2-6} -alkynyl groups respectively, especially C_{1-4} -alkyl, C_{2-4} -alkenyl and C_{2-4} -alkynyl groups, respectively and the halogen atoms fluoro, chloro, bromo or iodo atoms.

~~186~~

20 ~~24~~. (Amended) Epothilone derivative according to claim ~~24~~, wherein the aromatic and hetero aromatic, respectively, is provided with 1, 2 or 3 substituents and the hetero aromatic is provided with 1, 2 or more hetero atoms.

Sub
28
copy
13

C